

## Examples of QF Rings without Nakayama Automorphism and H-Rings without Self-Duality

Kazutoshi Koike

metadata, citation and similar papers at [core.ac.uk](http://core.ac.uk)

E-mail: [koike@c.oshima-k.ac.jp](mailto:koike@c.oshima-k.ac.jp)

Communicated by Kent R. Fuller

Received July 25, 2000

J. Kado and K. Oshiro (*J. Algebra* **211** (1999), 384–408) proved the equivalence of (A) the existence of a Nakayama isomorphism for every basic left H-ring (Harada ring) and (B) the existence of a Nakayama automorphism for every basic QF ring. In this paper, by using examples of J. Kraemer (“Characterizations of the Existence of (Quasi-) Self-Duality for Complete Tensor Rings,” *Algebra Berichte* 56, Verlag Reinhard Fischer, Munich, 1987), we give examples of QF rings without Nakayama automorphism and H-rings without self-duality. © 2001 Academic Press

*Key Words:* QF rings; H-rings; self-duality; Nakayama automorphism.

### 1. INTRODUCTION

By investigating the structure of H-rings deeply, Kado and Oshiro [5] proved that the following are equivalent:

- (A) the existence of a Nakayama isomorphism for every basic left H-ring,
- (B) the existence of a Nakayama automorphism for every basic QF ring,
- (C) the existence of self-duality for every basic left H-ring.

On the other hand, Kraemer [6] constructed an example of a QF ring that does not have a weakly symmetric self-duality. In Section 2 we shall point out that Kraemer’s example is an example of a QF ring without Nakayama automorphism (i.e., a counterexample of (B)) and give some other examples of QF rings without Nakayama automorphism. In Section 3, by giving

a necessary and sufficient condition for a special type of H-rings to have a self-duality (Theorem 3.1 and Corollary 3.1), we shall give concrete examples of H-rings without self-duality (i.e., a counterexample of (C)).

Throughout this paper, all rings have identity, all modules are unitary, and all homomorphisms are operated on the opposite side of scalars. For a module  $M$ , we denote, by  $E(M)$ ,  $J(M)$ ,  $S(M)$ , and  $T(M)$ , the injective hull, the radical, the socle, and the top of  $M$  (i.e., the factor module by its radical), respectively. For a ring  $R$ , we denote by  $R\text{-Mod}$  the category of left  $R$ -modules. For a left (resp. right)  $R$ -module  $X$  and a subset  $A$  of  $R$ , we denote by  $r_X(A)$  (resp.  $l_X(A)$ ) the annihilator of  $A$  in  $X$ .

## 2. QF RINGS WITHOUT NAKAYAMA AUTOMORPHISM

In this section, by presenting the example of Kraemer [6], we give examples of QF rings without Nakayama automorphism.

We begin by recalling terminologies of Morita duality. We say that a bimodule  ${}_R U_S$  defines a *Morita duality* if  ${}_R U_S$  is faithfully balanced and is an injective cogenerator on both sides. In particular, if  $R = S$ , we say that  ${}_R U_R$  defines a *self-duality*. A ring  $R$  is said to have a *left Morita duality* if there exists a bimodule  ${}_R U_S$  that defines a Morita duality. In this case, we say that  $R$  is *left Morita dual* to  $S$ . (See [9].)

Let  ${}_R U_R$  define a self-duality. We say that  ${}_R U_R$  defines a *weakly symmetric self-duality* if  $T({}_R Re) \cong {}_R \text{Hom}_R(T(eR_R), U_R)$  for every primitive idempotent  $e$  of  $R$ . Note that the condition of weakly symmetric self-duality is left-right symmetric. Since  $\text{Hom}_R(T(eR_R), U_R) \cong S({}_R Ue)$ ,  ${}_R U_R$  defines a weakly symmetric self-duality if and only if  $T({}_R Re) \cong S({}_R Ue)$  for every primitive idempotent  $e$  of  $R$ . A ring  $R$  is said to have a weakly symmetric self-duality if there exists a bimodule that defines a weakly symmetric self-duality. (See [6, p. 12].)

*Remark 2.1.* (1) As is well known, every artin algebra  $R$  has a weakly symmetric self-duality. Let  $K$  be the center of  $R$  and let  $E = E(T(K))$  be the minimal injective cogenerator for  $K\text{-Mod}$ . Then  $U = \text{Hom}_K(R, E)$ , which becomes an  $(R, R)$ -bimodule naturally, defines a weakly symmetric self-duality.

(2) A QF ring  $R$  is said to be weakly symmetric if the regular bimodule  ${}_R R_R$  defines a weakly symmetric self-duality.

According to [5], we give the following definition. Let  $R$  be a basic semiperfect ring with left Morita duality. Let  $\{e_1, e_2, \dots, e_n\}$  be a complete set of orthogonal primitive idempotents for  $R$  and let  $S = \text{End}_R(\bigoplus_{i=1}^n E(T(Re_i)))$  be the endomorphism ring of a minimal injective cogenerator for  $R\text{-mod}$ . Let  $f_i$  be the idempotent of  $S$  corresponding to

the projection  $\bigoplus_{i=1}^n E(T(Re_i)) \rightarrow E(T(Re_i))$ . Then we call a ring isomorphism  $\tau: R \rightarrow S$  a *Nakayama isomorphism* if  $\tau(e_i) = f_i$  for each  $i = 1, 2, \dots, n$ . By [7, p. 42], the existence of a Nakayama isomorphism does not depend on the choice of the complete set  $\{e_1, e_2, \dots, e_n\}$  of orthogonal primitive idempotents. (See [5, Remark on p. 387].)

The following proposition shows that the existence of a weakly symmetric self-duality coincides with that of a Nakayama isomorphism.

**PROPOSITION 2.1** ([4, Proposition 3.1]). *Let  $R$  be a basic semiperfect ring with left Morita duality and let  ${}_R U$  be a minimal injective cogenerator for  $R\text{-Mod}$ . Then the following conditions are equivalent:*

- (1)  *$R$  has a weakly symmetric self-duality.*
- (2)  *$R$  has a Nakayama isomorphism.*
- (3) *There exists a ring isomorphism  $\tau: R \rightarrow \text{End}({}_R U)$  such that  $U\tau(e) \cong E(T(Re))$  for every primitive idempotent  $e$  of  $R$ .*

*Proof.* (1)  $\Leftrightarrow$  (3) is [4, Proposition 3.1] and (2)  $\Leftrightarrow$  (3) is clear. ■

We recall the concepts of Nakayama permutations and Nakayama automorphisms for QF rings. Let  ${}_R U_S$  be a bimodule that defines a Morita duality. Then by [1, Lemma 30.2] we have  $S({}_R U) = S(U_S)$ . So we simply denote this module by  $S(U)$ . Let  $\{e_1, e_2, \dots, e_n\}$  and  $\{f_1, f_2, \dots, f_n\}$  be basic sets of orthogonal primitive idempotents for  $R$  and  $S$ , respectively. Then there exists a permutation  $\pi$  on  $\{1, 2, \dots, n\}$  such that  $S(e_i U) \cong T(f_{\pi(i)} S)$  and  $S(U f_{\pi(i)}) \cong T(Re_i)$  for each  $i = 1, 2, \dots, n$ . In particular, when  $R$  is a QF ring with a basic set  $\{e_1, e_2, \dots, e_n\}$  of orthogonal primitive idempotents, there exists a permutation  $\sigma$  on  $\{e_1, e_2, \dots, e_n\}$  such that  $S(e_i R) \cong T(\sigma(e_i) R)$  and  $S(R \sigma(e_i)) \cong T(Re_i)$  for each  $i = 1, 2, \dots, n$ . We call this permutation a *Nakayama permutation* of  $R$ . A ring automorphism  $\tau$  of  $R$  is said to be a *Nakayama automorphism* of  $R$  if the restriction  $\tau$  on  $\{e_1, e_2, \dots, e_n\}$  gives a Nakayama permutation of  $R$ . By Proposition 2.1, a basic QF ring  $R$  has a Nakayama automorphism if and only if  $R$  has a weakly symmetric self-duality.

We now give a condition to have a Nakayama automorphism for QF rings of a special type. For the sake of convenience, we denote by  $[i]$  the least positive residue of an integer  $i$  modulo a positive integer  $m$ . In the proof of the next proposition (and that of Theorem 3.1), we shall use the useful concept of *i-pair*. Let  $e, f$  be primitive idempotents of  $R$ . A pair  $(eR, Rf)$  is called an *i-pair* if  $S(eR) \cong T(fR)$  and  $S(Rf) \cong T(Re)$  (see [2]). As is well known (e.g., see [1, Theorem 31.3]), when  $R$  is a one-sided artinian ring, for a primitive idempotent  $e$  of  $R$ ,  $eR_R$  is injective if and only if there exists a primitive idempotent  $f$  of  $R$  such that  $(eR, Rf)$  is an *i-pair*. We shall use these results freely. Furthermore, we use the following

terminology. Let  $\alpha: A \rightarrow A'$  and  $\beta: B \rightarrow B'$  be ring homomorphisms and let  ${}_A M_B$  and  ${}_{A'} M'_{B'}$  be bimodules. An additive homomorphism  $\phi: M \rightarrow M'$  is said to be  $(\alpha, \beta)$ -semilinear if  $\phi(amb) = \alpha(a)\phi(m)\beta(b)$  for all  $a \in A$ ,  $b \in B$ , and  $m \in M$ .

PROPOSITION 2.2. *Let  $A_1, A_2, \dots, A_m$  ( $m \geq 2$ ) be basic artinian rings and let  ${}_{A_1}U_{1A_2}, {}_{A_2}U_{2A_3}, \dots, {}_{A_m}U_{mA_1}$  be bimodules, each of which defines a Morita duality. Let*

$$R = \begin{pmatrix} A_1 & U_1 & 0 & \cdots & 0 & 0 \\ 0 & A_2 & U_2 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & A_{m-1} & U_{m-1} \\ U_m & 0 & 0 & \cdots & 0 & A_m \end{pmatrix}$$

and define a ring structure on  $R$  by using ordinary matrix operations and by  $U_i U_j = 0$  for  $1 \leq i, j \leq m$ . Then

(1)  $R$  is a QF ring.

(2)  $R$  has a Nakayama automorphism if and only if for each  $i = 1, 2, \dots, m$ , there exists a ring isomorphism  $\tau_i: A_i \rightarrow A_{[i+1]}$  such that  $T(A_i e) \cong S(U_i \tau_i(e))$  for every primitive idempotent  $e$  of  $A_i$ , and there exists a  $(\tau_i, \tau_{[i+1]})$ -semilinear isomorphism  $\phi_i: U_i \rightarrow U_{[i+1]}$ .

*Proof.* (1) By assumption, all of  $A_i$  have complete sets consisting of the same number ( $n$ , say) of orthogonal primitive idempotents. Thus, for each  $i = 1, 2, \dots, m$ , there exists a complete set  $\{e_{ij} | 1 \leq j \leq n\}$  of orthogonal primitive idempotents for  $A_i$ . Then, since  ${}_{A_i}U_{iA_{[i+1]}}$  defines a Morita duality, there exists a permutation  $\pi_i$  on  $\{1, 2, \dots, n\}$  such that  $S(e_{ij}U_i) \cong T(e_{[i+1], \pi_i(j)}A_{[i+1]})$  and  $S(U_i e_{[i+1], \pi_i(j)}) \cong T(A_i e_{ij})$  for  $1 \leq j \leq n$ . For each  $i = 1, 2, \dots, m$  and  $j = 1, 2, \dots, n$ , let  $f_{ij}$  be the idempotent of  $R$  such that its  $(i, i)$ -entry is  $e_{ij}$  and all other entries are zero, and let  $f_i = \sum_{j=1}^n f_{ij}$  be the idempotent of  $R$  such that its  $(i, i)$ -entry is  $1_{A_i}$  and all other entries are zero. Then  $\{f_{ij} | 1 \leq i \leq m, 1 \leq j \leq n\}$  is a complete set of orthogonal primitive idempotents for  $R$ . By the definition of  $R$ , it is routine to check that  $S(f_{ij}R) \cong T(f_{[i+1], \pi_i(j)}R)$  and  $S(Rf_{[i+1], \pi_i(j)}) \cong T(Rf_{ij})$  for  $1 \leq i \leq m$  and  $1 \leq j \leq n$ . Thus  $(f_{ij}R, Rf_{[i+1], \pi_i(j)})$  become i-pairs and hence  $R$  is a QF ring.

(2)  $(\Rightarrow)$  For the setting above, the Nakayama permutation of  $R$  is given by  $\sigma: f_{ij} \mapsto f_{[i+1], \pi_i(j)}$ . Assume that  $R$  has a Nakayama automorphism  $\tau$ . Then  $\tau(f_{ij}) = f_{[i+1], \pi_i(j)}$  for  $1 \leq i \leq m$  and  $1 \leq j \leq n$ . Since

$$\tau(f_i) = \tau\left(\sum_{j=1}^n f_{ij}\right) = \sum_{j=1}^n \tau(f_{ij}) = \sum_{j=1}^n f_{[i+1], \pi_i(j)} = f_{[i+1]},$$

it follows from  $A_i \cong f_i R f_i$  and  $U_i \cong f_i R f_{[i+1]}$  that  $\tau$  induces ring isomorphisms  $\tau_i: A_i \rightarrow A_{[i+1]}$  and additive isomorphisms  $\phi_i: U_i \rightarrow U_{[i+1]}$ . It is clear that  $\tau_i(e_{ij}) = e_{[i+1], \pi_i(j)}$ , so  $T(A_i e_{ij}) \cong S(U_i e_{[i+1], \pi_i(j)}) = S(U_i \tau_i(e_{ij}))$ . Thus  $T(A_i e) \cong S(U_i \tau_i(e))$  for every primitive idempotent  $e$  of  $A_i$ . By the definitions of  $\tau_i$  and  $\phi_i$ , we see that  $\phi_i$  is  $(\tau_i, \tau_{[i+1]})$ -semilinear.

( $\Leftarrow$ ) Assume that  $\tau_i$  and  $\phi_i$  are given. We define  $\tau: R \rightarrow R$  via

$$\tau \begin{pmatrix} a_1 & u_1 & \cdots & 0 \\ 0 & a_2 & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & u_{m-1} \\ u_m & 0 & \cdots & a_m \end{pmatrix} = \begin{pmatrix} \tau_m(a_m) & \phi_m(u_m) & \cdots & 0 \\ 0 & \tau_1(a_1) & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & \phi_{m-2}(u_{m-2}) \\ \phi_{m-1}(u_{m-1}) & 0 & \cdots & \tau_{m-1}(a_{m-1}) \end{pmatrix}.$$

Then it is routine to check that  $\tau$  is a ring automorphism of  $R$  such that  $T(Rf) \cong S(R\tau(f))$  for every primitive idempotent  $f$  of  $R$ . Therefore by Proposition 2.1,  $R$  has a Nakayama automorphism. ■

*Remark 2.2.* In the setting of Proposition 2.2, let  $A = A_1 \times A_2 \times \cdots \times A_m$  be the product of rings and let  $U = U_1 \oplus U_2 \oplus \cdots \oplus U_m$  be the direct sum of  $\mathbb{Z}$ -modules. Then  $U$  becomes an  $(A, A)$ -bimodule naturally and  ${}_A U_A$  defines a self-duality. Thus by [9, Theorem 10.7] the trivial extension of  $A$  by  $U$  is a QF ring. Indeed,  $R$  is isomorphic to the trivial extension.

We now present Kraemer's example. To state this, we need the following notation. Let  $C$  and  $D$  be rings. If  ${}_C M_D$  is a bimodule, then let  $M_1 = {}_C M_D$  and for  $i = 2, 3, \dots$ , define inductively

$$M_i = \begin{cases} {}_C \text{Hom}_C({}_D M_{i-1}, {}_C C_C)_D & \text{if } i \text{ is odd,} \\ {}_D \text{Hom}_D({}_C M_{i-1}, {}_D D_D)_C & \text{if } i \text{ is even.} \end{cases}$$

EXAMPLE 2.1. By the results of [8] and [3], there exists an extension  $C > D$  of division rings satisfying the following conditions (see [6, Theorems 6.1 and 6.2]).

- (1)  $\dim({}_D C) = 2$  and  $\dim(C_D) = 3$ .
- (2) There exist ring isomorphisms  $\lambda: D \rightarrow C$  and  $\mu: C \rightarrow D$ .
- (3) There exists a  $(\lambda, \mu)$ -semilinear isomorphism  $\phi: {}_D C_{6C} \rightarrow {}_C C_{1D}$ .
- (4)  $(a_1, a_2, a_3, a_4, a_5) = (3, 1, 2, 2, 1)$  and  $(b_1, b_2, b_3, b_4, b_5) = (1, 3, 1, 2, 2)$ ,

where  $a_i$  and  $b_i$  denote the right and left dimensions of  $C_i$ , respectively. Then by [6, Lemma 6.3],

(5) The map  $\psi: {}_C C_{7D} \rightarrow {}_D C_{2C}$  defined by  $\psi(c_7)(c_1) = \mu(c_7(\phi^{-1}(c_1)))$  ( $c_7 \in C_7$ ,  $c_1 \in C_1$ ) is a  $(\mu, \lambda)$ -semilinear isomorphism.

Let

$$A_i = \begin{pmatrix} C & C_i \\ 0 & D \end{pmatrix} \quad (i \text{ is odd}) \quad \text{and} \quad A_i = \begin{pmatrix} D & C_i \\ 0 & C \end{pmatrix} \quad (i \text{ is even}),$$

the upper triangular matrix rings, and let

$$U_i = \begin{pmatrix} D & 0 \\ C_i & C \end{pmatrix} \quad (i \text{ is odd}) \quad \text{and} \quad U_i = \begin{pmatrix} C & 0 \\ C_i & D \end{pmatrix} \quad (i \text{ is even}).$$

Then by [9, Corollary 10.3],  $U_{i+1}$  becomes  $(A_{i+2}, A_i)$ -bimodules, which defines a Morita duality. By using  $\lambda$ ,  $\mu$ ,  $\phi$ , and  $\psi$ , we see that  $A_6 \cong A_1$  and  $A_7 \cong A_2$  as rings. Therefore we regard  $U_5$  as an  $(A_1, A_4)$ -bimodule that defines a Morita duality. Similarly, we regard  $U_1$  as an  $(A_2, A_5)$ -bimodule that defines a Morita duality. Now let

$$R = \begin{pmatrix} A_5 & U_4 & 0 & 0 & 0 \\ 0 & A_3 & U_2 & 0 & 0 \\ 0 & 0 & A_1 & U_5 & 0 \\ 0 & 0 & 0 & A_4 & U_3 \\ U_1 & 0 & 0 & 0 & A_2 \end{pmatrix}.$$

Using the dimensions  $a_i$  and  $b_i$  of  $C_i$ , we see that all  $A_i$  ( $1 \leq i \leq 5$ ) are pairwise non-isomorphic. Therefore by Proposition 2.2  $R$  is a QF ring that does not have a Nakayama automorphism.  $R$  is the ring of [6, Remark 6.6], which was given as an example of a QF ring without weakly symmetric self-duality.

EXAMPLE 2.2. Let  $A$  be an artinian ring with self-duality but without weakly symmetric self-duality and let  ${}_A U_A$  be a bimodule that defines a self-duality. (See Examples 2.1 and 2.3.) Then by Proposition 2.2 the ring

$$R = \begin{pmatrix} A & U & 0 & \cdots & 0 & 0 \\ 0 & A & U & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & A & U \\ U & 0 & 0 & \cdots & 0 & A \end{pmatrix}$$

is a QF ring without Nakayama automorphism.

We give another example of a QF ring without Nakayama automorphism.

EXAMPLE 2.3. With the same setting as in Example 2.1, let

$$A = \begin{pmatrix} C & C_1 & 0 & 0 & 0 \\ 0 & D & C_2 & 0 & 0 \\ 0 & 0 & C & C_3 & 0 \\ 0 & 0 & 0 & D & C_4 \\ C_5 & 0 & 0 & 0 & C \end{pmatrix}, \quad B = \begin{pmatrix} C & C_3 & 0 & 0 & 0 \\ 0 & D & C_4 & 0 & 0 \\ 0 & 0 & C & C_5 & 0 \\ 0 & 0 & 0 & D & C_6 \\ C_7 & 0 & 0 & 0 & C \end{pmatrix},$$

$$U = \begin{pmatrix} C & 0 & 0 & 0 & C_6 \\ C_2 & D & 0 & 0 & 0 \\ 0 & C_3 & C & 0 & 0 \\ 0 & 0 & C_4 & D & 0 \\ 0 & 0 & 0 & C_5 & C \end{pmatrix},$$

where we regard  $C_5$  in  $A$ ,  $C_6$  in  $U$ , and  $C_7$  in  $B$  as  $(C, C)$ -bimodules by using the ring isomorphism  $\lambda^{-1}: C \rightarrow D$ . Then by [6, Theorem 6.4]  $U$  becomes a  $(B, A)$ -bimodule that defines a Morita duality and there exists a ring isomorphism  $B \rightarrow A$  given by

$$\begin{pmatrix} a & b & 0 & 0 & 0 \\ 0 & c & d & 0 & 0 \\ 0 & 0 & e & f & 0 \\ 0 & 0 & 0 & g & h \\ j & 0 & 0 & 0 & i \end{pmatrix} \mapsto \begin{pmatrix} \lambda(g) & \phi(h) & 0 & 0 & 0 \\ 0 & \mu(i) & \psi(j) & 0 & 0 \\ 0 & 0 & a & b & 0 \\ 0 & 0 & 0 & c & d \\ f & 0 & 0 & 0 & e \end{pmatrix}.$$

Therefore we can regard  $U$  as an  $(A, A)$ -bimodule, which defines a self-duality. Let  $R$  be the trivial extension of  $A$  by  $U$ . Then  $R$  is a QF ring by [9, Theorem 10.7]. Let  $e'_i$  be the  $(i, i)$ -matrix unit of  $A$  and let  $e_i$  be the idempotent of  $R$  corresponding to  $e'_i$ . We denote by  $i$  a composition factor that is isomorphic to  $T(e_i R)$  or to  $T(R e_i)$  ( $i = 1, 2, \dots, 5$ ). By using the dimensions  $a_i$  and  $b_i$  of  $C_i$ , we see that the composition diagrams of the Loewy factors of the indecomposable projective modules of  $R_R$  and  ${}_R R$  are the following ones.

$R_R$

$$\begin{array}{ccccc} 1 & 2 & 3 & 4 & 5 \\ 2 & 2 & 2 & 3 & 3 \\ 4 & 5 & 1 & 2 & 2 \end{array}$$

${}_R R$ 

1	2	3	4	5
4 4 4 5 5	5 1	1 1 2 2 2	2 2 3	3 4 4
3	4	5	1	2

Thus the Nakayama permutation of  $R$  is  $e_i \mapsto e_{[i+3]}$  and is cyclic. However, the lengths of  $e_1 R$  and  $e_4 R$  are different. Therefore there does not exist an automorphism  $\tau$  of  $R$  such that  $\tau(e_1) = e_4$  and hence  $R$  does not have a Nakayama automorphism.

### 3. H-RINGS WITHOUT SELF-DUALITY

In this section, by using the examples of QF rings without a Nakayama automorphism in the previous section and by giving a necessary and sufficient condition for a special type of H-rings to have a self-duality, we provide concrete examples of H-rings without self-duality.

A left artinian ring  $R$  is said to be a *left H-ring* (*Harada ring*) if there exists a basic set  $\{e_{ij} | 1 \leq i \leq m, 1 \leq j \leq n(i)\}$  of orthogonal primitive idempotents for  $R$  satisfying the following conditions:

- (1)  $e_{i1} R_R$  is injective for each  $i = 1, 2, \dots, m$ .
- (2)  $e_{ij} R \cong J(e_{i, j-1} R_R)$  for each  $j = 2, 3, \dots, n(i)$ .

Right H-rings are defined similarly (cf. [5]). If  $R$  is a left H-ring, then by [5, Proposition 3.2] a minimal injective cogenerator  $\bigoplus_{i,j} E(T(Re_{ij}))$  is finitely generated. Therefore we note that  $R$  is left Morita dual to  $\text{End}_R(\bigoplus_{i,j} E(T(Re_{ij})))$  by [1, Theorem 30.4].

As we mentioned in the Introduction, in [5] it is shown that (A), (B), and (C) are equivalent. Therefore, since (A) and (B) do not hold from the examples in Section 2, there must exist a left H-ring without self-duality.

The next theorem, which can be seen as a special case of [5, Proposition 3.3], gives rise to concrete examples of such rings. To state this, we use the following notation. Let  $R$  be a ring with a complete set  $I$  of orthogonal primitive idempotents. For a non-empty subset  $K$  of  $I$ , we put  $e_K = \sum_{e \in K} e$ . For a nonzero idempotent  $e$  of  $R$ , we define

$$R_e = \begin{pmatrix} eRe & eRe & eR(1-e) \\ J(eRe) & eRe & eR(1-e) \\ (1-e)Re & (1-e)Re & (1-e)R(1-e) \end{pmatrix}.$$

Then  $R_e$  is a ring by usual matrix operations. We note that



$$\begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} R_e \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \cong \begin{pmatrix} eRe & eR(1-e) \\ (1-e)Re & (1-e)R(1-e) \end{pmatrix} \cong R.$$

**THEOREM 3.1.** *Let  $R$  be a basic QF ring and let  $I$  be a complete set of orthogonal primitive idempotents for  $R$  with the Nakayama permutation  $\sigma$  on  $I$ . For a non-empty subset  $K$  of  $I$ , let  $e = e_K$  and  $e' = e_{\sigma(K)}$ . Then*

(1)  $R_e$  is a left and right H-ring.

(2)  $R_e$  is left Morita dual to  $R_{e'}$ .

(3)  $R_e$  has a self-duality if and only if there exists a ring automorphism  $\tau$  of  $R$  such that  $\tau(e) = e'$ .

*Proof.* (1) Let  $A = eRe$ ,  $B = (1-e)R(1-e)$ ,  $U = eR(1-e)$ ,  $V = (1-e)Re$ , and  $\Lambda = R_e$ . Then

$$R = \begin{pmatrix} A & U \\ V & B \end{pmatrix}, \quad \Lambda = \begin{pmatrix} A & A & U \\ J(A) & A & U \\ V & V & B \end{pmatrix}.$$

Let  $I_A$  and  $I_B$  be complete sets of orthogonal primitive idempotents for  $A$  and  $B$ , respectively. We regard  $I$  as the disjoint union  $I_A \cup I_B$  and the Nakayama permutation  $\sigma$  as a permutation on  $I_A \cup I_B$ . For  $X, Y \in \{A, B\}$ , let  $I_{XY} = \{f \in I_X \mid \sigma(f) \in I_Y\}$ . Then  $I_A = I_{AA} \cup I_{AB}$  and  $I_B = I_{BA} \cup I_{BB}$ . For  $f \in I_A$  and  $g \in I_B$ , let

$$f^{(1)} = \begin{pmatrix} f & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad f^{(2)} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & f & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad g^{(3)} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & g \end{pmatrix}.$$

Since  $R$  is a basic QF ring, we have

$$J(R) = \begin{pmatrix} J(A) & U \\ V & J(B) \end{pmatrix}, \quad J(\Lambda) = \begin{pmatrix} J(A) & A & U \\ J(A) & J(A) & U \\ V & V & J(B) \end{pmatrix}.$$

Therefore, letting

$$S(R) = l_R(J(R)) = r_R(J(R)) = \begin{pmatrix} S_{11} & S_{12} \\ S_{21} & S_{22} \end{pmatrix},$$

since  $S(\Lambda_A) = l_\Lambda(J(\Lambda))$  and  $S_s(\Lambda) = r_\Lambda(J(\Lambda))$ , we have

$$S(\Lambda_\Lambda) = \begin{pmatrix} 0 & S_{11} & S_{12} \\ 0 & S_{11} & S_{12} \\ 0 & S_{21} & S_{22} \end{pmatrix}, \quad S({}_\Lambda\Lambda) = \begin{pmatrix} S_{11} & S_{11} & S_{12} \\ 0 & 0 & 0 \\ S_{21} & S_{21} & S_{22} \end{pmatrix}.$$

By using these representations of socles, it follows that  $(f^{(1)}\Lambda, \Lambda g^{(2)})$  ( $f \in I_{AA}$ ),  $(f^{(1)}\Lambda, \Lambda g^{(3)})$  ( $f \in I_{AB}$ ),  $(f^{(3)}\Lambda, \Lambda g^{(2)})$  ( $f \in I_{BA}$ ), and  $(f^{(3)}\Lambda, \Lambda g^{(3)})$  ( $f \in I_{BB}$ ) are i-pairs, where  $g = \sigma(f)$ . Therefore,  $f^{(1)}\Lambda_\Lambda$  ( $f \in I_A$ ),  $f^{(3)}\Lambda_\Lambda$  ( $f \in I_B$ ),  ${}_\Lambda\Lambda f^{(2)}$  ( $f \in I_A$ ), and  ${}_\Lambda\Lambda f^{(3)}$  ( $f \in I_B$ ) are injective. Also it is clear that  $f^{(2)}\Lambda_\Lambda \cong J(f^{(1)}\Lambda_\Lambda)$  and  ${}_\Lambda\Lambda f^{(1)} \cong J({}_\Lambda\Lambda f^{(2)})$  for  $f \in I_A$ . Therefore  $\Lambda$  is a left and right H-ring.

(2) We first compute a minimal injective cogenerator for  $\Lambda$ -Mod. For  $f \in I$ , let  $g = \sigma(f)$ . It follows from the i-pairs of the proof of (1) above and [5, Proposition 3.2] that

$$\begin{aligned} E(T(\Lambda f^{(1)})) &\cong \begin{cases} \Lambda g^{(2)} & (f \in I_{AA}), \\ \Lambda g^{(3)} & (f \in I_{AB}), \end{cases} \\ E(T(\Lambda f^{(2)})) &\cong \begin{cases} \Lambda g^{(2)}/S(\Lambda g^{(2)}) & (f \in I_{AA}), \\ \Lambda g^{(3)}/S(\Lambda g^{(3)}) & (f \in I_{AB}), \end{cases} \\ E(T(\Lambda f^{(3)})) &\cong \begin{cases} \Lambda g^{(2)} & (f \in I_{BA}), \\ \Lambda g^{(3)} & (f \in I_{BB}). \end{cases} \end{aligned}$$

For  $X, Y \in \{A, B\}$ , let  $f_{XY} = \sum_{f \in I_{XY}} f$  and  $g_{XY} = \sum_{f \in I_{XY}} \sigma(f)$ . Note that  $f_{XY} \in X$  and  $g_{XY} \in Y$ . Furthermore, let  $g_A = g_{AA}^{(2)} + g_{AB}^{(3)}$  and  $g_B = g_{BA}^{(2)} + g_{BB}^{(3)}$ , and let  $E^{(1)} = E(T(\Lambda 1_A^{(1)}))$ ,  $E^{(2)} = E(T(\Lambda 1_A^{(2)}))$ , and  $E^{(3)} = E(T(\Lambda 1_B^{(3)}))$ . Then from the isomorphisms above we have  $E^{(1)} \cong \Lambda g_A$ ,  $E^{(2)} \cong \Lambda g_A/S(\Lambda g_A)$ , and  $E^{(3)} \cong \Lambda g_B$ . Since  $\{1_A^{(1)}, 1_A^{(2)}, 1_B^{(3)}\}$  is a complete set of orthogonal idempotents for  $\Lambda$ ,  $E^{(1)} \oplus E^{(2)} \oplus E^{(3)}$  is a minimal injective cogenerator for  $\Lambda$ -Mod.

From  $1_A = g_{AA} + g_{BA}$ ,  $1_B = g_{AB} + g_{BB}$ , and the definition of  $e'$ , we now note that

$$e' = \begin{pmatrix} g_{AA} & 0 \\ 0 & g_{AB} \end{pmatrix}, \quad 1 - e' = \begin{pmatrix} g_{BA} & 0 \\ 0 & g_{BB} \end{pmatrix} \quad \text{in } R = \begin{pmatrix} A & U \\ V & B \end{pmatrix}.$$

Then by using the isomorphisms of  $E^{(i)}$  above and the fact that  $r_R(S(R)) = J(R)$ , it is routine to check that

$$\begin{aligned} &\text{End}_\Lambda(E^{(1)} \oplus E^{(2)} \oplus E^{(3)}) \\ &\cong \begin{pmatrix} \text{Hom}_\Lambda(E^{(1)}, E^{(1)}) & \text{Hom}_\Lambda(E^{(1)}, E^{(2)}) & \text{Hom}_\Lambda(E^{(1)}, E^{(3)}) \\ \text{Hom}_\Lambda(E^{(2)}, E^{(1)}) & \text{Hom}_\Lambda(E^{(2)}, E^{(2)}) & \text{Hom}_\Lambda(E^{(2)}, E^{(3)}) \\ \text{Hom}_\Lambda(E^{(3)}, E^{(1)}) & \text{Hom}_\Lambda(E^{(3)}, E^{(2)}) & \text{Hom}_\Lambda(E^{(3)}, E^{(3)}) \end{pmatrix} \end{aligned}$$

$$\cong \begin{pmatrix} e'Re' & e'Re' & e'R(1-e') \\ J(e'Re') & e'Re' & e'R(1-e') \\ (1-e')Re' & (1-e')Re' & (1-e')R(1-e') \end{pmatrix} = R_{e'}.$$

(3) ( $\Rightarrow$ ) Assume that  $\Lambda$  has a self-duality. Let  $A' = e'Re'$ ,  $B = (1-e')R(1-e')$ ,  $U' = e'R(1-e')$ ,  $V' = (1-e')Re'$ , and  $\Lambda' = R_{e'}$ . Then

$$\Lambda' = \begin{pmatrix} A' & A' & U' \\ J(A') & A' & U' \\ V' & V' & B' \end{pmatrix}.$$

By (2) there exists a ring isomorphism  $\rho: \Lambda \rightarrow \Lambda'$ . Let  $I_{A'}$  and  $I_{B'}$  be complete sets of orthogonal primitive idempotents for  $A'$  and  $B'$ , respectively, and let

$$L = \{f^{(1)}|f \in I_A\} \cup \{f^{(2)}|f \in I_A\} \cup \{f^{(3)}|f \in I_B\},$$

$$L' = \{f'^{(1)}|f' \in I_{A'}\} \cup \{f'^{(2)}|f' \in I_{A'}\} \cup \{f'^{(3)}|f' \in I_{B'}\}.$$

Then  $L$  and  $L'$  are complete sets of orthogonal primitive idempotents for  $\Lambda$  and  $\Lambda'$ , respectively. By [7, p. 42] we may assume that  $\rho(L) = L'$ . Since the injectivities of indecomposable projective  $\Lambda'$ -modules are similar to  $\Lambda$  in the proof of (1) and  $\rho$  preserves injective one-sided ideals, we must have  $\rho(\{f^{(1)}|f \in I_A\}) = \{f'^{(1)}|f' \in I_{A'}\}$  and  $\rho(\{f^{(2)}|f \in I_A\}) = \{f'^{(2)}|f' \in I_{A'}\}$ . Thus,  $\rho(1_A^{(1)}) = 1_{A'}^{(1)}$  and  $\rho(1_A^{(2)}) = 1_{A'}^{(2)}$ . Therefore  $\rho$  induces a ring isomorphism

$$\rho': \begin{pmatrix} A & U \\ V & B \end{pmatrix} \rightarrow \begin{pmatrix} A' & U' \\ V' & B' \end{pmatrix} \quad \text{such that } \rho' \begin{pmatrix} 1_A & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 1_{A'} & 0 \\ 0 & 0 \end{pmatrix}.$$

Then it follows from the definitions of these two rings that  $\rho'$  induces a ring automorphism  $\tau$  of  $R$  such that  $\tau(e) = e'$ .

( $\Leftarrow$ ) Assume that there exists a ring automorphism  $\tau$  of  $R$  such that  $\tau(e) = e'$ . Then we define  $\rho: R_e \rightarrow R_{e'}$  via  $\rho((r_{ij})_{i,j}) = (\tau(r_{ij}))_{i,j}$ . Since  $\tau(e) = e'$ ,  $\rho$  is well defined and is a ring isomorphism. Therefore by (2)  $R_e$  has a self-duality. ■

*Remark 3.1.* In the theorem above, the ring  $R_e$  does not have a self-duality; that is,  $R_e \not\cong R_{e'}$  in general. However, since the Nakayama permutation  $\sigma$  has a finite order, by iterating to take left Morita dual rings, we return to the first ring  $R_e$ .

The following corollary is a special case of Theorem 3.1.

**COROLLARY 3.1.** *Let  $R$  be a basic QF ring and let  $I$  be a basic set of*

orthogonal primitive idempotents for  $R$  with the Nakayama permutation  $\sigma$  on  $I$ . Assume that there exists a disjoint union  $I = I_1 \cup U_2 \cup \cdots \cup I_m$  such that  $\sigma(I_i) = I_{[i+1]}$  for each  $i$ . Let  $e_i = e_{I_i}$ ,  $R_{ij} = e_i R e_j$ , and

$$\Lambda = \begin{pmatrix} R_{11} & R_{11} & R_{12} & \cdots & R_{1m} \\ J(R_{11}) & R_{11} & R_{12} & \cdots & R_{1m} \\ R_{21} & R_{21} & R_{22} & \cdots & R_{2m} \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ R_{m1} & R_{m1} & R_{m2} & \cdots & R_{mm} \end{pmatrix}.$$

Then

- (1)  $\Lambda$  is a left and right H-ring.
- (2)  $\Lambda$  is left Morita dual to

$$\Gamma = \begin{pmatrix} R_{22} & R_{22} & R_{23} & \cdots & R_{2m} & R_{21} \\ J(R_{22}) & R_{22} & R_{23} & \cdots & R_{2m} & R_{21} \\ R_{32} & R_{32} & R_{33} & \cdots & R_{3m} & R_{31} \\ \vdots & \vdots & \vdots & & \vdots & \vdots \\ R_{m2} & R_{m2} & R_{m3} & \cdots & R_{mm} & R_{m1} \\ R_{12} & R_{12} & R_{13} & \cdots & R_{1m} & R_{11} \end{pmatrix}.$$

- (3)  $\Lambda$  has a self-duality if and only if there exists a ring automorphism  $\tau$  of  $R$  such that  $\tau(e_1) = e_2$ .

*Proof.* (1) and (3). By noting that  $\Lambda = R_{e_1}$  and  $\Gamma = R_{e_2}$  and with the definition of  $e_i$ , these are special cases of Theorem 3.1 (1) and (3).

(2) Let  $f_0$  be the idempotent of  $\Lambda$  such that its  $(1, 1)$ -entry is  $1_{A_1}$  and all other entries are zero, and for  $i = 1, 2, \dots, m$ , let  $f_i$  be the idempotent of  $\Lambda$  such that its  $(i+1, i+1)$ -entry is  $1_{A_i}$  and all other entries are zero. As well as the proof of Theorem 3.1, it follows that  $E(T(\Lambda f_0)) \cong \Lambda f_2$ ,  $E(T(\Lambda f_1)) \cong \Lambda f_2 / S(\Lambda f_2)$ ,  $E(T(\Lambda f_2)) \cong \Lambda f_3, \dots, E(T(\Lambda f_{m-1})) \cong \Lambda f_m$ , and  $E(T(\Lambda f_m)) \cong \Lambda f_1$ . By using these isomorphisms, it is routine to check that the endomorphism ring  $\Gamma$  of a minimal injective cogenerator  $\bigoplus_{i=0}^m E(T(\Lambda f_i))$  has the desired form. ■

Concluding this paper, we give two concrete examples of H-rings without self-duality.

EXAMPLE 3.1. Let  $A_i$  and  $U_i$  ( $i = 1, 2, \dots, 5$ ) be the same as in Example 2.1. Therefore  ${}_{A_{[i+2]}}U_{[i+1]A_i}$  defines a Morita duality and all

$A_1, A_2, \dots, A_5$  are pairwise non-isomorphic. Let

$$\Lambda = \begin{pmatrix} A_5 & A_5 & U_4 & 0 & 0 & 0 \\ J(A_5) & A_5 & U_4 & 0 & 0 & 0 \\ 0 & 0 & A_3 & U_2 & 0 & 0 \\ 0 & 0 & 0 & A_1 & U_5 & 0 \\ 0 & 0 & 0 & 0 & A_4 & U_3 \\ U_1 & U_1 & 0 & 0 & 0 & A_2 \end{pmatrix}.$$

Since  $A_5$  and  $A_3$  are not isomorphic,  $\Lambda$  is a left and right H-ring without self-duality by Corollary 3.1.

Since each  $A_i$  is a  $2 \times 2$  matrix ring,  $\Lambda$  is a  $12 \times 12$  matrix ring. Let  $e_i$  be the  $(i, i)$ -matrix unit and we denote by  $i$  a composition factor that is isomorphic to  $T(e_i \Lambda)$  or to  $T(\Lambda e_i)$  ( $i = 1, 2, \dots, 12$ ). Then the composition diagrams of the Loewy factors of the indecomposable projective modules of  $\Lambda_\Lambda$  and  ${}_\Lambda \Lambda$  are the following.

$\Lambda_\Lambda$

1	2	3	4	5	6
3	4	2	5 5	6 6	7
2	5 5	4	6	7	8
4	6	5			
5					
7	8	9	10	11	12
8 8 8	9	10 10	11 11	12	1 1 1
9	10	11	12	1	3 3 3
				3	2
					4

${}_\Lambda \Lambda$

1	2	3	4	5	6
12	3 3	1	2	4 4	5
11	1 1	12	3 3	2 2	4
	12	11	1 1	3	2
			12	1	
7	8	9	10	11	12
6 6 6	7	8 8	9 9	10	11 11 11
5	6	7	8	9	10

EXAMPLE 3.2. Let  $R$  be the QF ring without a Nakayama automorphism in Example 2.3 and let  $e_1, e_2, \dots, e_5$  be the primitive idempotents for  $R$ . The Nakayama permutation of  $R$  is  $e_i \mapsto e_{[i+3]}$  and is cyclic. Let

$$\Lambda = R_{e_1} = \begin{pmatrix} e_1 R e_1 & e_1 R e_1 & e_1 R(1 - e_1) \\ J(e_1 R e_1) & e_1 R e_1 & e_1 R(1 - e_1) \\ (1 - e_1) R e_1 & (1 - e_1) R e_1 & (1 - e_1) R(1 - e_1) \end{pmatrix}.$$

Then  $\Lambda$  has six isomorphism classes of the simple modules. By example 2.3 the composition lengths of  $e_1 \Lambda$  and  $e_4 \Lambda$  are different. Therefore by Theorem 3.1  $\Lambda$  is a left and right H-ring without self-duality.

## REFERENCES

1. F. W. Anderson and K. R. Fuller, "Rings and Categories of Modules," 2nd ed., Springer-Verlag, Berlin/Heidelberg/New York, 1992.
2. Y. Baba and K. Oshiro, On a theorem of Fuller, *J. Algebra* **154** (1993), 86–94.
3. P. Dowbor, C. M. Ringel, and D. Simson, "Hereditary Artinian Rings and Finite Representation Type, Lecture Notes in Mathematics, Vol. 832, pp. 232–241, Springer-Verlag, Berlin/Heidelberg/New York, 1980.
4. J. K. Haack, Self-duality and serial rings, *J. Algebra* **59** (1979), 345–363.
5. J. Kado and K. Oshiro, Self-duality and Harada rings, *J. Algebra* **211** (1999), 384–408.
6. J. Kraemer, "Characterizations of the Existence of (Quasi-) Self-Duality for Complete Tensor Rings," *Algebra Berichte* 56, Verlag Reinhard Fischer, Munich, 1987.
7. C. Faith, "Algebra II: Ring Theory," Springer-Verlag, Berlin/Heidelberg/New York, 1976.
8. A. H. Schofield, "Representations of Rings over Skew Fields," Cambridge Univ. Press, Cambridge, U.K., 1985.
9. W. Xue, "Rings with Morita Duality," Lecture Notes in Mathematics, Vol. 1523, Springer-Verlag, Berlin/Heidelberg/New York, 1992.